

# **hPSMA**

**Nomenclature** C57BL/6Smoc-*Folh1*<sup>em1(hPSMA-Wpre-PA)Smoc</sup>

**Cat. NO.** NM-HU-200240

**Strain State** Embryo cryopreservation

### **Gene Summary**

| Gene Symbol<br>Folh1 | Synonyms       | GCP2; mopsm       |
|----------------------|----------------|-------------------|
|                      | NCBI ID        | <u>53320</u>      |
|                      | MGI ID         | 1858193           |
|                      | Ensembl ID     | ENSMUSG0000001773 |
|                      | Human Ortholog | FOLH1B            |

## **Model Description**

The endogenous mice Folh1 gene was replaced by human PSMA gene.

\*Literature published using this strain should indicate: hPSMA mice (Cat. NO. NM-HU-200240) were purchased from Shanghai Model Organisms Center, Inc..

#### **Validation Data**








Fig1. Detection of PSMA expression in kidney by RT-PCR. Wild type: only one band at 294 bp with primers F1/R1(mPsma); Heterozygous: one band at 294 bp with primers F1/R1(mPsma) and one band at 169 bp with primers F2/R2(hPSMA); Abbr.. M, DNA marker; HO, homozygous; HE, heterozygous; WT, wild type.




Fig2. Detection of hPSMA expression in prostate in hPSMA KI mice.




Fig3. Analysis of human PSMA expression in the MC38-hPSMA cell line and the transplanted tumor of the PSMA humanized mice by FACS. Most of the MC38-hPSMA cells express human PSMA (A,B); the transplanted tumor formed by MC38-hPSMA cell line expresses human PSMA on the



CD45- cells (C).(Completed in collaboration with CrownBio)

### Mean Tumor Volume ± SEM




Fig4. Subcutaneous tumor formation experiment of the MC38-hPSMA cells in the PSMA humanized mice. The tumor volume was measured every 5 days after injection of MC38-hPSMA cells (n=10). The previous studies have shown that wild type mice would spontaneously reject the syngeneic tumors expressing an introduced human tumor antigen-hPSMA. While the MC38-hPSMA cells can form tumors subcutaneously in the humanized PSMA mice, indicating that the PSMA humanized mouse model can be used to transplant the MC38-hPSMA cell line. (Completed in collaboration with CrownBio)




Fig5. Detection of human PSMA expression in Prostate by IHC.

Note. The human PSMA antibody cross-reacted with mouse PSMA and human PSMA.



| Abbr. HO, homozyg | jous; WT, wild type. |  |  |
|-------------------|----------------------|--|--|
|                   |                      |  |  |
|                   |                      |  |  |
|                   |                      |  |  |
|                   |                      |  |  |
|                   |                      |  |  |